

TB/HIV PBMEF Indicators

Note on TB/HIV indicators: For activities supported by PEPFAR, PEPFAR indicators should be used as written in the Monitoring, Evaluation, and Reporting (MER) Indicator Reference Guide and reported through appropriate PEPFAR channels. These indicators are offered in the PBMEF as a recommended way to analyze these data points and for use outside of PEPFAR programming. To avoid confusion, the PBMEF describes PEPFAR indicator equivalencies where they exist and adopts PEPFAR naming conventions rather than the PBMEF conventions.

Table of Contents:

TB/HIV PBMEF Indicators	1
PCT_TX_CURR_SCRN: Percentage of PLHIV on ART screened for TB disease	2
TX_TB_D_MWRD: Number of PLHIV screened for TB with mWRD	3
TX_TB_D_CXR: Number of PLHIV screened for TB with chest X-ray	4
TX_TB_D_LAM: Number of PLHIV screened for TB with LF-LAM	4
TX_TB_D_CRP: Number of PLHIV screened for TB with CRP	5
PCT_TX_TB_SCRN_POS: Percentage of PLHIV on ART, screened positive for TB disease	6
TX_TB_SCRN_POS_MWRD: Number of PLHIV screened TB positive with mWRD	6
TX_TB_SCRN_POS_CXR: Number of PLHIV screened TB positive with chest X-ray	7
TX_TB_SCRN_POS_LAM: Number of PLHIV screened TB positive with LF-LAM	8
TX_TB_SCRN_POS_CRP: Number of PLHIV screened positive for TB with CRP	9
PCT_TX_TB_TST: Percentage of PLHIV on ART with a specimen sent	9
PCT_TX_TB_TST_POS: Percentage of PLHIV on ART with a positive TB result returned	10
PCT_TX_DRTB_TST_POS: Percentage of PLHIV on ART with a positive DR-TB result returned	11
PCT_TX_TB: Percentage of PLHIV who enrolled on TB treatment	11
PCT_TX_DRTB: Percentage of PLHIV who enrolled on DR-TB treatment	12
PCT_TBHIV_OUT: TB/HIV coinfecting treatment outcomes	13
TBHIV_TSR: Treatment success rate among PLHIV	13
PCT_TBHIV_DIED: TB/HIV coinfecting treatment outcome: Died during treatment	13
PCT_TBHIV_FAIL: TB/HIV coinfecting treatment outcome: Treatment failed	14
PCT_TBHIV_LTFU: TB/HIV coinfecting treatment outcome: LTFU	15
PCT_TBHIV_NE: TB/HIV coinfecting treatment outcome: Not evaluated	15
DRTBHIV_TSR: DR-TB/HIV coinfection treatment success rate	16
PCT_DRTBHIV_DIED: DR-TB/HIV coinfection treatment outcome: Died during treatment	17
PCT_DRTBHIV_FAIL: DR-TB/HIV coinfecting treatment outcome: Treatment failed	17
PCT_DRTBHIV_LTFU: DR-TB/HIV coinfecting treatment outcome: LTFU	18
PCT_DRTBHIV_NE: DR-TB/HIV coinfecting treatment outcome: Not evaluated	18
PCT_TX_TB_SCRN_NEG: Percentage of PLHIV on ART, screened negative for TB	19
TB_PREV_D: Number of TPT initiations among PLHIV	20

PCT_TX_CURR_TPT: Percentage PLHIV enrolled on TPT	20
PCT_TB_PREV: Percentage of PLHIV completed TPT	20
PCT_TB_PREV_LTFU: Percentage of PLHIV on TPT: LTFU	21
PCT_TB_PREV_TB: Percentage of PLHIV on TPT developed TB during TPT	22
PCT_TB_PREV_ADR: Percentage of PLHIV on TPT, and TPT interrupted due to adverse drug reaction (ADR)	22
PCT_TB_PREV_AST: Percentage of PLHIV on TPT, with baseline AST/ALT tests	23
PCT_TB_STAT: Percentage of people with TB with known HIV status	24
PCT_DRTB_STAT: Percentage of people with DR-TB with known HIV status	24
PCT_TB_STAT_POS: Percentage of people with TB recorded as HIV-positive	25
PCT_DRTB_STAT_POS: Percentage of people with DR-TB recorded as HIV-positive	26
PCT_TB_ART: Percentage of HIV-positive people with TB on ART	27
PCT_DRTB_ART: Percentage of HIV-positive people with DR-TB on ART	28
PCT_TB_ART_AHD: PLHIV diagnosed with TB who have <200 CD4 baseline	28
PCT_TB_STAT_POS_VL: PLHIV diagnosed with TB with baseline viral load results	29

The color of the indicator reference sheet indicates whether the indicator is calculated or not calculated, and whether it could be considered a disaggregate of another indicator:

Calculated indicators:

Indicator name	<i>Ex: PCT_TB_ART: Percentage TB clients who start or continue ART</i>
----------------	--

Non calculated indicators:

Indicator name	<i>Ex: TB_ART_N: Number of TB clients who start or continue ART</i>
----------------	---

Calculated indicators that could be reported as a disaggregate of another indicator:

Indicator name	<i>Ex: PCT_DRTB_ART: Percentage DR-TB clients who start or continue ART</i>
----------------	---

Non calculated indicators that could be reported as a disaggregate of another indicator:

Indicator name	<i>Ex: DRTB_ART_N: Number of DR-TB clients who start or continue ART</i>
----------------	--

Indicator name	PCT_TX_CURR_SCRN: Percentage of PLHIV on ART screened for TB disease <i>Previously [TH-1]</i>
Definition	Percentage of people living with HIV (PLHIV) enrolled on antiretroviral therapy (ART) who were screened at least once for TB, among all PLHIV enrolled on ART during the same reporting period. Note: This indicator is equivalent to the PEPFAR indicators <i>TX_TB denominator /</i>

	<p><i>TX_CURR</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who were screened at least once for TB during the reporting period	<p><i>PBMEF data element: TX_TB_D*</i></p> <p><i>*Note this is a PEPFAR indicator.</i></p>
Denominator	Number of PLHIV enrolled on ART during the reporting period	<p><i>PBMEF data element: TX_CURR*</i></p> <p><i>*Note this is a PEPFAR indicator.</i></p>
Category	Reach	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	Age, sex, screening type (mWRD, chest X-ray, CRP, LF-LAM, symptom screen alone), HIV treatment status (new vs previously enrolled on treatment for 6mo or longer)	

Indicator name	TX_TB_D_MWRD: Number of PLHIV screened for TB with mWRD	
Definition	<p>Number of PLHIV screened for TB with molecular WHO-recommended rapid diagnostic (mWRD).</p> <p>Note: This indicator is equivalent to the mWRD disaggregate for screening type in the denominator of the PEPFAR indicator <i>TX_TB</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV screened for TB with mWRD	<p><i>PBMEF data element: TX_TB_D_mWRD or TX_TB_D disaggregate*</i></p> <p><i>*Note this is a PEPFAR indicator disaggregate.</i></p>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	

Unit of measure	Number of people
Disaggregate by	This indicator could be recorded as a disaggregate of TX_TB_D.

Indicator name	TX_TB_D_CXR: Number of PLHIV screened for TB with chest X-ray	
Definition	<p>Number of people living with HIV (PLHIV) screened for TB with chest X-ray.</p> <p>Note: This indicator is equivalent to the CXR disaggregate for screening type in the denominator of the PEPFAR indicator <i>TX_TB</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV screened for TB with chest X-ray	<p><i>PBMEF data element: TX_TB_D_CXR or TX_TB_D disaggregate*</i></p> <p><i>*Note this is a PEPFAR indicator disaggregate.</i></p>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	
Unit of measure	Number of people	
Disaggregate by	This indicator could be recorded as a disaggregate of TX_TB_D.	

Indicator name	TX_TB_D_LAM: Number of PLHIV screened for TB with LF-LAM	
Definition	<p>Number of people living with HIV (PLHIV) screened for TB with LF-LAM</p> <p>LF-LAM is included as a recommended TB test for people living with HIV (PLHIV). LF-LAM is not recommended to confirm TB in all populations and notably should not be used in outpatient settings for adults, adolescents, and children without symptoms of TB or in those with a CD4 count > 200 cells/mm³. At the time of this publication, Alere Determine™ TB LAM Ag is the only commercially available LF- LAM test. Full guidance on the use of LF-LAM can be found at: www.who.int/publications/i/item/9789241550604.</p>	

	Note: This indicator could be a disaggregate for the denominator of the PEPFAR indicator TX_TB, however PEPFAR does not include an option for LF-LAM reporting. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.	
Numerator	Number of PLHIV screened for TB with LF-LAM	<i>PBMEF data element: TX_TB_D_LAM or TX_TB_D disaggregate*</i> <i>*Note this disaggregate is not included in the PEPFAR indicator.</i>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	
Unit of measure	Number of people	
Disaggregate by	This indicator could be recorded as a disaggregate of TX_TB_D.	

Indicator name	TX_TB_D_CRP: Number of PLHIV screened for TB with CRP	
Definition	Number of people living with HIV (PLHIV) screened for TB with C-reactive protein (CRP). Note: This indicator could be a disaggregate for the denominator of the PEPFAR indicator TX_TB, however PEPFAR does not include an option for CRP reporting. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.	
Numerator	Number of PLHIV screened for TB with CRP	<i>PBMEF data element: TX_TB_D_CRP or TX_TB_D disaggregate*</i> <i>*Note this disaggregate is not included in the PEPFAR indicator.</i>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	

Unit of measure	Number of people
Disaggregate by	This indicator could be recorded as a disaggregate of TX_TB_D.

Indicator name	PCT_TX_TB_SCRN_POS: Percentage of PLHIV on ART, screened positive for TB Previously [TH-2]	
Definition	<p>Percentage of people living with HIV (PLHIV) enrolled on ART who were screened positive for TB, among all PLHIV enrolled on ART who were screened for TB during the same reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicators <i>TX_TB denominator screen positive (TX_TB POS) / TX_TB denominator</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who screened positive for TB during the reporting period	<i>PBMEF data element: TX_TB_SCRN_POS*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Denominator	Number of PLHIV enrolled on ART who screened at least once for TB during the reporting period	<i>PBMEF data element: TX_TB_D*</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Reach	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	Age, sex, screening type (mWRD, chest X-ray, symptom screen alone), HIV treatment status (new vs previously enrolled on treatment for 6mo or longer)	

Indicator name	TX_TB_SCRN_POS_MWRD: Number of PLHIV screened TB positive with mWRD	
Definition	<p>Number of people living with HIV (PLHIV) screened positive for TB with molecular WHO-recommended rapid diagnostic (mWRD).</p> <p>Note: This indicator could be a further disaggregation for the screen positive disaggregate in the PEPFAR indicator <i>TX_TB</i>, however PEPFAR does not collect this data. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR</p>	

	programming.	
Numerator	Number of PLHIV screened positive for TB with mWRD	<i>PBMEF data element: TX_TB_SCRN_POS_MWRD or TX_TB_SCRN_POS disaggregate*</i> <i>*Note this disaggregate is not included in the PEPFAR indicator.</i>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	
Unit of measure	Number of people	
Disaggregate by	This indicator could be reported as a disaggregate of TX_TB_SCRN_POS.	

Indicator name	TX_TB_SCRN_POS_CXR: Number of PLHIV screened TB positive with chest X-ray	
Definition	<p>Number of people living with HIV (PLHIV) screened positive for TB with chest X-ray.</p> <p>Note: This indicator could be a further disaggregation for the screen positive disaggregate in the PEPFAR indicator TX_TB, however PEPFAR does not collect this data. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV screened positive for TB with chest X-ray	<i>PBMEF data element: TX_TB_SCRN_POS_CXR or TX_TB_SCRN_POS disaggregate*</i> <i>*Note this disaggregate is not included in the PEPFAR indicator.</i>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	
Unit of measure	Number of people	

Disaggregate by	This indicator could be reported as a disaggregate of TX_TB_SCRN_POS.
------------------------	---

Indicator name	TX_TB_SCRN_POS_LAM: Number of PLHIV screened TB positive with LF-LAM	
Definition	<p>Number of people living with HIV (PLHIV) screened positive for TB with LF-LAM</p> <p>LF-LAM is included as a recommended TB test for people living with HIV (PLHIV). LF- LAM is not recommended to confirm TB in all populations and notably should not be used in outpatient settings for adults, adolescents, and children without symptoms of TB or in those with a CD4 count > 200 cells/mm3. At the time of this publication, Alere Determine™ TB LAM Ag is the only commercially available LF- LAM test. Full guidance on the use of LF-LAM can be found at:www.who.int/publications/i/item/9789241550604.</p> <p>Note: This indicator could be a further disaggregation for the screen positive disaggregate in the PEPFAR indicator <i>TX_TB</i>, however PEPFAR does not collect this data. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV screened positive for TB with LF-LAM	<p><i>PBMEF data element:</i> <i>TX_TB_SCRN_POS_LAM or</i> <i>TX_TB_SCRN_POS disaggregate*</i></p> <p><i>*Note this disaggregate is not included in the PEPFAR indicator.</i></p>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	
Unit of measure	Number of people	
Disaggregate by	This indicator could be reported as a disaggregate of TX_TB_SCRN_POS.	

Indicator name	TX_TB_SCRN_POS_CRP: Number of PLHIV screened positive for TB with CRP	
Definition	<p>Number of people living with HIV (PLHIV) screened positive for TB with C-reactive protein (CRP).</p> <p>Note: This indicator could be a further disaggregation for the screen positive disaggregate in the PEPFAR indicator <i>TX_TB</i>, however PEPFAR does not collect this data. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive</p>	

	PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.	
Numerator	Number of PLHIV screened positive for TB with CRP	<i>PBMEF data element: TX_TB_SCRN_POS_CRP or TX_TB_SCRN_POS disaggregate*</i> <i>*Note this disaggregate is not included in the PEPFAR indicator.</i>
Denominator	N/A	N/A
Category	Prevent	
Indicator type	Output	
PBMEF level	Extended	
Unit of measure	Number of people	
Disaggregate by	This indicator could be reported as a disaggregate of TX_TB_SCRN_POS.	

Indicator name	PCT_TX_TB_TST: Percentage of PLHIV on ART with a specimen sent Previously [TH-3]	
Definition	<p>Percentage of people living with HIV (PLHIV) enrolled on ART who had a specimen sent for bacteriologic diagnosis of active TB disease, among PLHIV enrolled on ART who screened positive for TB during the reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicators <i>TX_TB denominator specimen sent / TX_TB denominator screen positive</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who had a specimen sent for diagnosis of active TB disease during the reporting period	<i>PBMEF data element: TX_TB_TST*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Denominator	Number of PLHIV enrolled on ART who screened positive for TB during the reporting period	<i>PBMEF data element: TX_TB_SCRN_POS*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Category	Reach	
Indicator type	Outcome; calculated	

PBMEF level	Extended
Unit of measure	Percent of people
Disaggregate by	

Indicator name	PCT_TX_TB_TST_POS: Percentage of PLHIV on ART with a positive TB result returned <i>Previously [TH-4]</i>	
Definition	<p>Percentage of people living with HIV (PLHIV) enrolled on ART who had a positive result returned for bacteriologic diagnosis of active TB disease, among PLHIV enrolled on ART who had a specimen sent for TB diagnosis during the reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicators <i>TX_TB denominator positive result returned / TX_TB denominator specimen sent</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who had a positive result returned for diagnosis of active TB disease during the reporting period	<i>PBMEF data element: TX_TB_TST_POS*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Denominator	Number of PLHIV enrolled on ART who had a specimen sent for diagnosis of active TB disease during the reporting period	<i>PBMEF data element: TX_TB_TST*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Category	Reach	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	DS-TB vs DR-TB	

Indicator name	PCT_TX_DRTB_TST_POS: Percentage of PLHIV on ART with a positive DR-TB result returned
Definition	Percentage of people living with HIV (PLHIV) enrolled on ART who had a positive result

	<p>returned for bacteriologic diagnosis of active DR-TB disease, among PLHIV enrolled on ART who had a specimen sent for TB diagnosis during the reporting period.</p> <p>Note: This indicator is a disaggregate of PEPFAR's TX_TB disaggregate for positive results returned, but is not included in PEPFAR reporting.</p>	
Numerator	Number of PLHIV enrolled on ART who had a positive result returned for diagnosis of active DR-TB disease during the reporting period	<i>PBMEF data element: TX_DRTB_TST_POS*</i> <i>*Note this disaggregate of TX_TB_D is not included in PEPFAR's indicator.</i>
Denominator	Number of PLHIV enrolled on ART who had a specimen sent for diagnosis of active TB disease during the reporting period	<i>PBMEF data element: TX_TB_TST*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Category	Reach	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator is a disaggregate of PCT_TX_TB_TST_POS.	

Indicator name	PCT_TX_TB: Percentage of PLHIV who enrolled on TB treatment <i>Previously [TH-5]</i>	
Definition	<p>Percentage of people living with HIV (PLHIV) enrolled on ART who were enrolled on TB treatment, among PLHIV who had a positive result returned for diagnosis of active TB disease during the reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicator TX_TB. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who were enrolled on TB treatment, during the reporting period	<i>PBMEF data element: TX_TB_N*</i> <i>*Note this is a PEPFAR indicator.</i>
Denominator	Number of PLHIV enrolled on ART who had a positive result returned for diagnosis of active TB disease during the reporting period	<i>PBMEF data element: TX_TB_TST_POS*</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Cure	
Indicator type	Outcome; calculated	

PBMEF level	Extended
Unit of measure	Percent of people
Disaggregate by	

Indicator name	PCT_TX_DRTB: Percentage of PLHIV who enrolled on DR-TB treatment <i>Previously [TH-17]</i>	
Definition	Percentage of PLHIV enrolled on ART who were enrolled on DR-TB treatment, among PLHIV who had a positive result returned for diagnosis of active DR-TB disease during the reporting period.	
Numerator	Number of PLHIV enrolled on ART who were enrolled on DR-TB treatment, during the reporting period	<i>PBMEF data element: TX_DRTB_N*</i> <i>*Note this disaggregate of TX_TB_N is not included in PEPFAR's indicator.</i>
Denominator	Number of PLHIV enrolled on ART who had a positive result returned for diagnosis of active DR-TB disease during the reporting period	<i>PBMEF data element: TX_DRTB_TST_POS*</i> <i>*Note this disaggregate of TX_TB_D is not included in PEPFAR's indicator.</i>
Category	Cure	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be recorded as a disaggregate of PCT_TX_TB.	

Indicator name	PCT_TBHIV_OUT: TB/HIV coinfectd treatment outcomes	
Definition	Percentage TB/HIV coinfectd people with a new episode of TB (new, recurrent ¹) who died during treatment, among people coinfectd with TB/HIV (all forms, new and recurrent) on treatment during the same reporting period.	
Numerator	Number of TB/HIV coinfectd people (all forms) with a new episode of TB (new, recurrent) who had a recorded treatment outcome, during the reporting period; this should be disaggregated by the following	<i>PBMEF data element: TBHIV_SUCC</i> <i>WHO database: tbhiv_succ</i> <i>PBMEF data element: TBHIV_DIED</i> <i>WHO database: tbhiv_died</i>

	outcomes: treatment success, died, LTFU, not evaluated	<i>PBMEF data element: TBHIV_FAIL</i> <i>WHO indicator: tbhiv_fail</i> <i>PBMEF data element: TBHIV_LOST</i> <i>WHO database: tbhiv_lost</i> <i>PBMEF data element: TBHIV_NA</i>
Denominator	Number of TB/HIV coinfecting people with a new episode of TB (new, recurrent) (all forms) on treatment during the reporting period (cohort size)	<i>PBMEF data element: TBHIV_COH</i> <i>WHO indicator: tbhiv_coh</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT or as a disaggregate of TX_DS_OUT.	

¹Previously “relapse”

Indicator name	TBHIV_TSR: Treatment success rate among PLHIV
Note:	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

Indicator name	PCT_TBHIV_DIED: TB/HIV coinfecting treatment outcome: Died during treatment <i>Previously [TH-20]</i>	
Definition	Percentage TB/HIV coinfecting people with a new episode of TB (new, recurrent ¹) who died during treatment, among people coinfecting with TB/HIV (all forms, new and recurrent) on treatment during the same reporting period.	
Numerator	Number of TB/HIV coinfecting people (all forms) with a new episode of TB (new, recurrent) who died during treatment, during the reporting period	<i>PBMEF data element: TBHIV_DIED</i> <i>WHO database: tbhiv_died</i>
Denominator	Number of TB/HIV coinfecting people with a new episode of TB (new, recurrent) (all forms) on treatment during the reporting	<i>PBMEF data element: TBHIV_COH</i> <i>WHO indicator: tbhiv_coh</i>

	period (cohort size)	
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT or as a disaggregate of TX_DS_OUT.	

¹Previously “relapse”

Indicator name	PCT_TBHIV_FAIL: TB/HIV coinfecting treatment outcome: Treatment failed <i>Previously [TH-21]</i>	
Definition	Percentage TB/HIV coinfecting people with a new episode of TB (new, recurrent ¹) whose treatment failed, among people coinfecting with TB/HIV (all forms, new and recurrent) on treatment during the same reporting period.	
Numerator	Number of TB/HIV coinfecting people (all forms) with a new episode of TB (new, recurrent) whose treatment failed, during treatment, during the reporting period	<i>PBMEF data element: TBHIV_FAIL WHO indicator: tbhiv_fail</i>
Denominator	Number of TB/HIV coinfecting people with a new episode of TB (new, recurrent) (all forms) on treatment during the reporting period (cohort size)	<i>PBMEF data element: TBHIV_COH WHO indicator: tbhiv_coh</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT or as a disaggregate of TX_DS_OUT.	

¹Previously “relapse”

Indicator name	PCT_TBHIV_LTFU: TB/HIV coinfectd treatment outcome: LTFU <i>Previously [TH-22]</i>	
Definition	Percentage TB/HIV coinfectd people with a new episode of TB (new, recurrent ¹) who were LTFU, among people coinfectd with TB/HIV (all forms, new and recurrent) on treatment during the same reporting period.	
Numerator	Number of TB/HIV coinfectd people (all forms) with a new episode of TB (new, recurrent) who were LTFU, during treatment, during the reporting period	<i>PBMEF data element: TBHIV_LOST WHO database: tbhiv_lost</i>
Denominator	Number of TB/HIV coinfectd people with a new episode of TB (new, recurrent) (all forms) on treatment during the reporting period (cohort size)	<i>PBMEF data element: TBHIV_COH WHO indicator: tbhiv_coh</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT or as a disaggregate of TX_DS_OUT.	

¹Previously “relapse”

Indicator name	PCT_TBHIV_NE: TB/HIV coinfectd treatment outcome: Not evaluated <i>Previously [TH-23]</i>	
Definition	Percentage TB/HIV coinfectd people with a new episode of TB (new, recurrent ¹) who were not evaluated, among people coinfectd with TB/HIV (all forms, new and recurrent) on treatment during the same reporting period.	
Numerator	Number of TB/HIV coinfectd people (all forms) with a new episode of TB (new, recurrent) who were not evaluated, during treatment, during the reporting period	<i>PBMEF data element: TBHIV_NE</i>
Denominator	Number of TB/HIV coinfectd people with a new episode of TB (new, recurrent) (all forms) on treatment during the reporting period (cohort size)	<i>PBMEF data element: TBHIV_COH WHO indicator: tbhiv_coh</i>
Category	Cure	

Indicator type	Outcome
PBMEF level	Extended
Unit of measure	Percent of people
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT or as a disaggregate of TX_DS_OUT.

¹Previously “relapse”

Indicator name	DRTBHIV_TSR: DR-TB/HIV coinfection treatment success rate <i>Previously [TH-24]</i>	
Definition	Percentage of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period and had a successful outcome (cured or treatment completed).	
Numerator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period, and had a successful outcome (cured or treatment completed).	<i>PBMEF data element: DRTBHIV_SUCC</i>
Denominator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period	<i>PBMEF data element: DRTBHIV_COH</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be reported as a disaggregate of PLHIV_TSR or DR_TSR.	

Indicator name	PCT_DRTBHIV_DIED: DR-TB/HIV coinfection treatment outcome: Died during treatment <i>Previously [TH-24]</i>
-----------------------	--

Definition	Percentage of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period and died during treatment.	
Numerator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period, and died during treatment	<i>PBMEF data element: DRTBHIV_DIED</i>
Denominator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period	<i>PBMEF data element: DRTBHIV_COH</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percent of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT, PCT_PLHIV_OUT_DIED, or TX_DR_OUT.	

Indicator name	PCT_DRTBHIV_FAIL: DR-TB/HIV coinfecting treatment outcome: Treatment failed Previously [TH-25]	
Definition	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period, but treatment failed.	
Numerator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period, but treatment failed	<i>PBMEF data element: DRTBHIV_FAIL</i>
Denominator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period	<i>PBMEF data element: DRTBHIV_COH</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT,	

	PCT_PLHIV_OUT_FAIL, or TX_DR_OUT.
--	-----------------------------------

Indicator name	PCT_DRTBHIV_LTFU: DR-TB/HIV coinfecting treatment outcome: LTFU <i>Previously [TH-26]</i>	
Definition	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on appropriate treatment during a specified period, but were LTFU.	
Numerator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on appropriate treatment during the reporting period, but were LTFU	<i>PBMEF data element: DRTBHIV_LTFU</i>
Denominator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period	<i>PBMEF data element: DRTBHIV_COH</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT, PCT_PLHIV_OUT_LTFU, or TX_DR_OUT.	

Indicator name	PCT_DRTBHIV_NE: DR-TB/HIV coinfecting treatment outcome: Not evaluated <i>Previously [TH-27]</i>	
Definition	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period, but not evaluated (not evaluated includes "transferred out," "still on treatment," and any other notified case where the treatment outcome has not been evaluated).	
Numerator	Number of DR-TB/HIV coinfecting people (RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period, but not evaluated (not evaluated includes "transferred out," "still on treatment," and any other notified case where the treatment outcome has not been evaluated)	<i>PBMEF data element: DRTBHIV_NE</i>
Denominator	Number of DR-TB/HIV coinfecting people	<i>PBMEF data element: DRTBHIV_COH</i>

	(RR/MDR and XDR) who were enrolled on DR-TB treatment during the reporting period	
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	This indicator could be reported as a disaggregate of PCT_PLHIV_OUT, PCT_PLHIV_OUT_NE, or TX_DR_OUT.	

Indicator name	PCT_TX_TB_SCRN_NEG: Percentage of PLHIV on ART, screened negative for TB <i>Previously [TH-6]</i>	
Definition	<p>Percentage of people living with HIV (PLHIV) enrolled on ART who screened negative for TB disease, among all PLHIV enrolled on ART who were screened for TB during the same reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicators <i>TX_TB denominator screen negative / TX_TB denominator</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who screened negative for TB disease during the reporting period	<i>PBMEF data element: TX_TB_SCRN_NEG*</i> <i>*Note this is a PEPFAR indicator disaggregate.</i>
Denominator	PLHIV enrolled on ART who were screened for TB disease during the reporting period	<i>PBMEF data element: TX_TB_D*</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Reach	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	This indicator could be recorded as a disaggregate of TX_TB or TX_TB_SCRN.	

Indicator name	TB_PREV_D: Number of TPT initiations among PLHIV <i>Previously [PT-8]</i>
Note:	No feedback requested on this indicator as it is an essential indicator and already has an associated PIRS.

Indicator name	PCT_TX_CURR_TPT: Percentage PLHIV enrolled on TPT <i>Previously [PT-8]</i>	
Definition	<p>Percentage of people living with HIV (PLHIV) enrolled on ART who were started on TPT during the reporting period, among all PLHIV enrolled on ART.</p> <p>Note: This indicator is equivalent to the PEPFAR indicators <i>TB_PREV D / TX_CURR</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of PLHIV enrolled on ART who were started on TPT during the reporting period	<i>PBMEF data element: TB_PREV_D*</i> <i>WHO indicator (2020 and beyond): hiv_all_tpt or hiv_new_tpt</i> <i>*Note: this is a PEPFAR indicator.</i>
Denominator	Number of PLHIV enrolled on ART during the reporting period	<i>PBMEF data element: TX_CURR*</i> <i>WHO indicator: hiv_reg_all, hiv_reg_new2</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, TPT regimen type	

Indicator name	PCT_TB_PREV: Percentage of PLHIV completed TPT <i>Previously [PT-12]</i>
Definition	Percentage of people living with HIV (PLHIV) enrolled on ART who were started on TPT and completed treatment, among all PLHIV enrolled on ART who were started on TPT during the reporting period.

	Note: This indicator is equivalent to the PEPFAR indicator <i>TB_PREV</i> ($\% \text{ completion} = \text{numerator} / \text{denominator}$). PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.	
Numerator	Number of PLHIV enrolled on ART who were started on TPT and completed treatment during the reporting period	<i>PBMEF data element: TB_PREV_N*</i> <i>*Note: this is a PEPFAR indicator.</i>
Denominator	Number of PLHIV enrolled on ART who were started on TPT during the reporting period	<i>PBMEF data element: TB_PREV_D*</i> <i>WHO indicator (2020 and beyond): hiv_all_tpt or hiv_new_tpt</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, TPT regimen type	

Indicator name	PCT_TB_PREV_LTFU: Percentage of PLHIV on TPT: LTFU <i>Previously [TH-9]</i>	
Definition	Percentage of people living with HIV (PLHIV) who were started on TPT but were LTFU. Note defining patients as LTFU may vary by regimen type and country guidelines.	
Numerator	Number of PLHIV who were started on TPT and were LTFU during the reporting period	<i>PBMEF data element: TB_PREV_LTFU*</i> <i>Note: TB_PREV is a PEPFAR indicator but there is no disaggregate for LTFU.</i>
Denominator	Number of PLHIV who were started on TPT during the specified reporting period	<i>PBMEF data element: TB_PREV_D*</i> <i>WHO indicator (2020 and beyond): hiv_all_tpt or hiv_new_tpt</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome	
PBMEF level	Extended	

Unit of measure	Percentage of people
Disaggregate by	Age, sex, TPT regimen type

Indicator name	PCT_TB_PREV_TB: Percentage of PLHIV on TPT developed TB during TPT Previously [TH-10]	
Definition	Percentage of people living with HIV (PLHIV) on TPT who developed TB during prevention therapy.	
Numerator	Total number of PLHIV who were started on TPT and developed TB during TPT, during the reporting period	<i>PBMEF data element: TB_PREV_TB*</i> <i>Note: TB_PREV is a PEPFAR indicator but there is no disaggregate for TB diagnosis.</i>
Denominator	Total number of PLHIV who were started on TPT during the specified reporting period	<i>PBMEF data element: TB_PREV_D*</i> <i>WHO indicator (2020 and beyond): hiv_all_tpt or hiv_new_tpt</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, TPT regimen type	

Indicator name	PCT_TB_PREV_ADR: Percentage of PLHIV on TPT, and TPT interrupted due to adverse drug reaction (ADR) Previously [TH-11]	
Definition	<p>Percentage of people living with HIV (PLHIV) whose TPT was interrupted due to ADR (adverse drug reaction).</p> <p>An ADR (often referred to as an “adverse event”) is any negative medical occurrence that presents in a person during TB preventive treatment with a World Health Organization (WHO) approved regimen that may or may not have a causal relationship with the prescribed treatment. More information on ADR and grading ADRs can be found here.</p>	
Numerator	Total number of PLHIV who began on TPT but whose treatment was interrupted due to development of ADR, during the reporting period	<i>PBMEF data element: TB_PREV_ADR*</i> <i>Note: TB_PREV is a PEPFAR indicator but there is no disaggregate for ADRs.</i>

Denominator	Total number of PLHIV who were started on TPT during the specified reporting period	<i>PBMEF data element: TB_PREV_D*</i> <i>WHO indicator (2020 and beyond): hiv_all_tpt or hiv_new_tpt</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, TPT regimen type	

Indicator name	PCT_TB_PREV_AST: Percentage of PLHIV on TPT, with baseline AST/ALT tests <i>Previously [TH-12]</i>	
Definition	Percentage of people living with HIV (PLHIV) on TB preventive treatment (TPT) who had a baseline alanine aminotransferase (ALT) and aspartate aminotransferase (AST) before initiation of prevention therapy.	
Numerator	Total number of PLHIV who had baseline AST/ALT tests before initiation of TPT, during the reporting period	<i>PBMEF data element: TB_PREV_AST*</i> <i>Note: TB_PREV is a PEPFAR indicator but there is no disaggregate for AST/ALT.</i>
Denominator	Total number of PLHIV who were started on TPT during the specified reporting period	<i>PBMEF data element: TB_PREV_D*</i> <i>WHO indicator (2020 and beyond): hiv_all_tpt or hiv_new_tpt</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, TPT regimen type	

Indicator name	PCT_TB_STAT: Percentage of people with TB with known HIV status <i>Previously [TH-13]</i>	
-----------------------	---	--

Definition	Percentage of people with new and relapse TB notified during the reporting period who were tested for HIV at the time of diagnosis or with known HIV status at the time of TB diagnosis, among all people with new and relapse TB (all forms) notified during the reporting period.	
	Note: This indicator is equivalent to the PEPFAR indicator <i>TB_STAT</i> . PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.	
Numerator	Number of people with new and relapse TB with a documented HIV status during the reporting period	<i>PBMEF data element: TB_STAT_N*</i> <i>WHO data element: newrel_hivtest</i> <i>*Note: this is a PEPFAR indicator.</i>
Denominator	Number of people with new and relapse TB (all forms) notified during the reporting period	<i>PBMEF data element: TB_STAT_D*</i> <i>WHO data element: c_newinc</i> <i>*Note: this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, DS-TB vs DR-TB	

Indicator name	PCT_DRTB_STAT: Percentage of people with DR-TB with known HIV status <i>Previously [TH-15]</i>	
Definition	Percentage of people with DR-TB with a documented HIV status, among all DR-TB patients notified during the reporting period.	
Numerator	Number of people with DR-TB who were tested for HIV at the time of diagnosis or with known positive or recently tested negative* HIV status at the time of DR-TB diagnosis during the reporting period. *Recently tested negative: people with DR-TB who have a recent negative test result and are not yet eligible for re-testing per country guidelines.	<i>PBMEF data element: TB_STAT disaggregate*</i> <i>*Note: TB_STAT is a PEPFAR indicator but this disaggregate is not included.</i>
Denominator	Number of people with laboratory-confirmed DR-TB notified during reporting year	<i>PBMEF data element: MDR_NOTIF + XDR_NOTIF</i> <i>WHO indicator: conf_rrmdr plus</i>

		<i>all_conf_xdr</i>
Category	Cure	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, HIV status for numerator (tested negative, tested positive, known positive, recent negative) This indicator is a disaggregate of PCT_TB_STAT.	

Indicator name	PCT_TB_STAT_POS: Percentage of people with TB recorded as HIV-positive Previously [TH-14]	
Definition	<p>Percentage of people with new and relapse TB recorded as HIV-positive, among all people with new and relapse TB (all forms) with a documented HIV status during the reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicator <i>TB_STAT_POS / TB_STAT numerator</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of people with new and relapse TB recorded as HIV-positive during the reporting period	<i>PBMEF data element: TB_STAT_POS*</i> <i>WHO data element: newrel_hivpos</i> <i>*Note this is a PEPFAR indicator.</i>
Denominator	Number of people with new and relapse TB with a documented HIV status, during the reporting period	<i>PBMEF data element: TB_STAT_N*</i> <i>WHO data element: newrel_hivtest</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Reach, cure, prevent, sustain, innovate	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex	

Indicator name	PCT_DRTB_STAT_POS: Percentage of people with DR-TB recorded as HIV-positive Previously [TH-16]	
Definition	Percentage of people with DR-TB documented as HIV-positive, among all people with DR-TB notified during the reporting period.	
Numerator	Number of people with DR-TB who were documented as HIV-positive during the reporting period	<i>PBMEF data element: DRTB_STAT_POS*</i> <i>*Note TB_STAT_POS is a PEPFAR indicator but the disaggregation for DR-TB is not included.</i>
Denominator	Number of people with DR-TB who had a documented HIV status during the reporting period	<i>PBMEF data element: DRTB_STAT_N*</i> <i>*Note TB_STAT_N is a PEPFAR indicator but the disaggregation for DR-TB is not included.</i>
Category	Cure	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex	

Indicator name	PCT_TB_ART: Percentage of HIV-positive people with TB on ART <i>Previously [TH-18]</i>	
Definition	<p>Percentage of HIV-positive people with TB started or continued on ART during the reporting period, among people with new and relapse TB recorded as HIV-positive during the reporting period.</p> <p>Note: This indicator is equivalent to the PEPFAR indicator <i>TB_ART</i>. PEPFAR indicators should be used as written in the MER Reference Guide for activities that receive PEPFAR funding and report through PEPFAR channels; this indicator is offered in the PBMEF as a recommended way to analyze this data and for use outside of PEPFAR programming.</p>	
Numerator	Number of HIV-positive people with TB started or continued on ART during the reporting period	<i>PBMEF data element: TB_ART_N*</i> <i>WHO data element: newrel_art, hiv_tbt_x_art</i> <i>*Note this is a PEPFAR indicator.</i>
Denominator	Number of people with new and relapse TB recorded as HIV-positive during the reporting period	<i>PBMEF data element: TB_STAT_POS*</i> <i>WHO data element: newrel_hivpos</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex, DS-TB vs DR-TB	

Indicator name	PCT_DRTB_ART: Percentage of HIV-positive people with DR-TB on ART <i>Previously [TH-19]</i>	
Definition	Percentage of people with DR-TB started or continued on ART during the reporting period, among people with DR-TB documented as HIV-positive during the reporting period.	
Numerator	Number of HIV-positive people with DR-TB started or continued on ART during the reporting period	<i>PBMEF data element: DRTB_ART*</i> <i>*Note TB_ART is a PEPFAR indicator but the disaggregation for DR-TB is not included.</i>
Denominator	Number of people with DR-TB documented as HIV-positive during the reporting period	<i>PBMEF data element: DRTB_STAT_POS*</i> <i>*Note TB_STAT_POS is a PEPFAR indicator but the disaggregation for DR-TB is not included.</i>
Category	Cure	
Indicator type	Outcome	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex	

Indicator name	PCT_TB_ART_AHD: PLHIV diagnosed with TB who have <200 CD4 baseline <i>Previously [TH-28]</i>	
Definition	Percentage of people living with HIV (PLHIV) diagnosed with TB who have <200 CD4 baseline starting TB treatment, among HIV-positive people with TB started or continued on ART during the reporting period.	
Numerator	Number of PLHIV diagnosed with TB who have <200 CD4 baseline starting treatment during the reporting period	<i>PBMEF data element: TB_ART_AHD*</i> <i>*Note TB_ART is a PEPFAR indicator but the disaggregate for AHD is not included.</i>
Denominator	Number of HIV-positive people with TB started or continued on ART during the reporting period	<i>PBMEF data element: TB_ART_N*</i> <i>WHO data element: newrel_art, hiv_tbt_x_art</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Prevent	
Indicator type	Outcome; calculated	
PBMEF level	Extended	

Unit of measure	Percentage of people
Disaggregate by	Age, sex

Indicator name	PCT_TB_STAT_POS_VL: PLHIV diagnosed with TB with baseline viral load results <i>Previously [TH-30]</i>	
Definition	Percentage of PLHIV diagnosed with TB with baseline viral load results available, among HIV-positive people with TB during the reporting period.	
Numerator	Number of PLHIV diagnosed with TB who has baseline viral load results available during the reporting period	<i>PBMEF data element: TB_STAT_POS_VL</i>
Denominator	Number of people with new and relapse TB recorded as HIV-positive during the reporting period	<i>PBMEF data element: TB_STAT_POS*</i> <i>WHO data element: newrel_hivpos</i> <i>*Note this is a PEPFAR indicator.</i>
Category	Cure	
Indicator type	Outcome; calculated	
PBMEF level	Extended	
Unit of measure	Percentage of people	
Disaggregate by	Age, sex	